

- variation in *Styracosaurus* and *Centrosaurus*. *Vertebrate Anatomy Morphology Palaeontology*, 8, 67-95. <https://doi.org/10.18435/vamp29361>.
- Campbell, J. A., Ryan, M. J., Schröder-Adams, C. J., Evans, D. C., & Holmes, R. B. (2018). New insights into Chasmosaurine (Dinosauria: Ceratopsidae) skulls from the Upper Cretaceous (Campanian) of Alberta, and an update on the distribution of accessory frill fenestrae in Chasmosaurinae. *PeerJ*, 6, e5194. <https://doi.org/10.7717/peerj.5194>.
- Chinsamy, A., & Tumarkin-Deratzian, A. (2009). Pathologic bone tissues in a vulture and a nonavian dinosaur: Implications for interpreting endosteal bone and radial fibrolamellar bone in fossil dinosaurs. *The Anatomical Record*, 292(9), 1478-1484. <https://doi.org/10.1002/ar.20991>.
- Currie, P. J., Langston W., & Tanke, D. H. (2008). *A new horned dinosaur from an Upper Cretaceous bone bed in Alberta*. Canadian Science Publishing. <https://doi.org/10.1139/9780660198194>.
- Dyer, A. D., LeBlanc, A. R. H., Doschak, M. R., & Currie, P. J. (2021). Taking a crack at the dome: Histopathology of a pachycephalosaurid (Dinosauria: Ornithischia) frontoparietal dome. *Biosis: Biological Systems*, 2(2), 248-270. <https://doi.org/10.37819/biosis.002.02.0101>.
- Farke, A. A., Wolff, E. D. S., & Tanke, D. H. (2009). Evidence of combat in *Triceratops*. *PLoS ONE*, 4(1), e4252. <https://doi.org/10.1371/journal.pone.0004252>.
- Farlow, J. O., & Dodson, P. (1975). The behavioral significance of frill and horn morphology in ceratopsian dinosaurs. *Evolution*, 29(2), 353-361.
- Frederickson, J. A., & Tumarkin-Deratzian, A. R. (2014). Craniofacial ontogeny in *Centrosaurus apertus*. *PeerJ*, 2, e252.
- Hatcher, J. B. (1907). *The Ceratopsia*, by John B. Hatcher, based on preliminary studies by Othniel C. Marsh, ed. and completed by Richard S. Lull. United States Geological Survey. Monographs; v. 49. Government Printing Office.
- Hieronimus, T. L., Witmer, L. M., Tanke, D. H., & Currie, P. J. (2009). The facial integument of centrosaurine ceratopsids: Morphological and histological correlates of novel skin structures. *The Anatomical Record*, 292(9), 1370-1396. <https://doi.org/10.1002/ar.20985>.
- Langston Jr., W. (1967). The thick-headed ceratopsian dinosaur *Pachyrhinosaurus* (Reptilia: Ornithischia), from the Edmonton Formation near Drumheller, Canada. *Canadian Journal of Earth Sciences*, 4(1), 171-186. <https://doi.org/10.1139/e67-008>.
- Moodie, R. L. (1923). *Paleopathology; An introduction to the study of ancient evidences of disease*. University of Illinois Press.
- Padian, K., & Lamm, E.-T. (2013). *Bone histology of fossil tetrapods: Advancing methods, analysis, and interpretation* (1st ed.). University of California Press.
- Peterson, J. E., Dischler, C., & Longrich, N. R. (2013). Distributions of cranial pathologies provide evidence for head-butting in dome-headed dinosaurs (Pachycephalosauridae). *PLoS ONE*, 8(7), e68620. <https://doi.org/10.1371/journal.pone.0068620>.
- Sampson, S. D., Ryan, M. J., & Tanke, D. H. (1997). Craniofacial ontogeny in centrosaurine dinosaurs (Ornithischia: Ceratopsidae): Taxonomic and behavioral implications. *Zoological Journal of the Linnean Society*, 121(3), 293-337. <https://doi.org/10.1111/j.1096-3642.1997.tb00340.x>.
- Stilson, K. T., Hopkins S. S. B., & Davis, E. B. (2016). Osteopathology in Rhinocerotidae from 50 million years to the present. *PLoS ONE*, 11(2), e0146221. <https://doi.org/10.1371/journal.pone.0146221>.
- Tanke, D. H., & Rothschild, B. M. (2010). Paleopathologies in Albertan ceratopsids and their behavioral significance. In M. J. Ryan, B. J. Chinnery-Allgeier, & D. A. Eberth (Eds.), *New perspectives on horned dinosaurs* (pp. 355-384). Indiana University Press.
- Tanke, D. H., & Farke, A. A. (2006). Bone resorption, bone lesions and extra cranial fenestrae in ceratopsid dinosaurs: A preliminary assessment. In K. Carpenter (Ed.), *Horns and beaks: Ceratopsian and ornithomimid dinosaurs* (pp. 319-347). Indiana University Press.
- Wolff, E. D. S., Salisbury, S. W., Horner, J. R., & Varricchio, D. J. (2009). Common avian infection plagued the tyrant dinosaurs. *PLoS ONE*, 4(9), e7288. <https://doi.org/10.1371/journal.pone.0007288>.
- Woodruff, D. C., Wolff, E. D. S., Wedel, M. J., Dennison, S., & Witmer, L. M. (2022). The first occurrence of an avian-style respiratory infection in a non-avian dinosaur. *Scientific Reports*, 12(1), 1954. <https://doi.org/10.1038/s41598-022-05761-3>.

### Technical Session 3: Terrestrial Ecosystems – Late Jurassic (Friday, June 9, 2023, 2:00 PM)

#### NEW MATERIAL OF *HAPLOCANTHOSAURUS* (HATCHER 1903) FROM THE DRY MESA DINOSAUR QUARRY AND A COMMENT ON SAUROPOD DIVERSITY

Curtice, Brian<sup>1</sup>, Wedel, Mathew<sup>2</sup>, Wilhite, Ray<sup>3</sup>, Boisvert, Colin<sup>4</sup>

<sup>1</sup>Fossil Crates, 1305 W. Auto Dr., Tempe, Arizona, USA 85284, [Brian@FossilCrates.com](mailto:Brian@FossilCrates.com); <sup>2</sup>College of Osteopathic Medicine of the Pacific, Western University of Health Sciences, Pomona, California, USA, [mathew.wedel@gmail.com](mailto:mathew.wedel@gmail.com), ORCID 0000-0001-6082-3103; <sup>3</sup>Auburn University, College of Veterinary Medicine, 1130 Wire Road, Auburn, Alabama, USA 36849, [DRW0004@Auburn.edu](mailto:DRW0004@Auburn.edu); <sup>4</sup>Brigham Young University, Department of Geosciences, S389 Eyring

Science Center (ESC), Provo, Utah, USA 84602,  
 cdboisvert1998@gmail.com

*Haplocanthosaurus priscus* (Hatcher, 1903), “simple-spined lizard,” was named from partial skeletons collected from the Marsh-Felch Quarry near Cañon City, Colorado. Another species, *H. delfsi*, was named from a partial skeleton collected from the Cleveland Delfs Quarry, also in Colorado (McIntosh and Williams, 1988). To these specimens, fragmentary remains from several other localities have been assigned to the genus (Foster, 2020), and a nearly complete skeleton that may represent a new species awaits publication (Bilbey et al., 2000).

Brigham Young University Museum of Paleontology houses five *Haplocanthosaurus* dorsal vertebrae (BYU 9194, 11506, 17530 (Figure 1), 17531, 17689) and a tibia (BYU 12865, Figure 1) from the Late Jurassic Dry Mesa Dinosaur Quarry (DMDQ). The dorsal vertebrae possess diagnostic *Haplocanthosaurus* characters, including tall neural arch peduncles, dorsally-angled transverse processes, and parapophyses placed well below the diapophyses even on posterior dorsal vertebrae (Figure 1). The right tibia is identified as *Haplocanthosaurus* based on the robust nature of the element and the general morphology of a greatly expanded distal end and small cnemial crest. The least breadth/length (LB/L) measurement of 0.21 is greater than any other known sauropod taxa from DMDQ and compares well with USNM 4275, which includes a left tibia assigned to *H. priscus* (LB/L 0.23). While the left tibia of USNM 4275 has an articulated astragalus on its distal end, the morphology matches well with BYU 12865.

The primary importance of the new specimens is stratigraphic rather than morphological. *Haplocanthosaurus* is not abundant at any locality in the Morrison Formation, and the Marsh-Felch Quarry in Colorado is the only quarry to definitively produce more than one individual of *Haplocanthosaurus* (CM 572, CM 879, and USNM 4275). Despite its numerical scarcity, *Haplocanthosaurus* persisted through much of the deposition of the Morrison Fm. Dorsal vertebrae referable to *Haplocanthosaurus* are present low in the Salt Wash Member of the Morrison Fm. in central Utah (MJW pers. obs., Figure 1), and other *Haplocanthosaurus* localities are in the upper Salt Wash, lower Brushy Basin, and upper Brushy Basin members of the Morrison Fm. (Turner & Peterson, 1999). The DMDQ *Haplocanthosaurus* material confirms the presence of this long-lived genus in the upper Brushy Basin (Britt, 1991). The *Haplocanthosaurus* elements reported herein from DMDQ do not help resolve the phylogenetic position of this enigmatic genus.

Hatcher (1903) considered *Haplocanthosaurus* to be “... the most generalized member of the Sauropoda yet

discovered in America”, with the species name *priscus* meaning “first.” Historically the phylogenetic relationships of *Haplocanthosaurus* have been uncertain. Various analyses have found it to be a derived eusauropod just outside of Neosauropoda, a basal macronarian alongside *Camarasaurus* and *Brachiosaurus*, or the most basal diplodocoid, with more recent analyses tending to support the diplodocoid hypothesis (see Foster & Wedel, 2014 for a brief review; Foster, 2020).

The Dry Mesa Dinosaur Quarry has produced the most diverse assemblage of sauropod dinosaurs in any one locality in the world. The quarry, opened in 1972, was excavated for the better part of two decades and, at over 120m long, it is one of the larger Morrison Formation localities. Unfortunately, the majority of the nearly 5,000 elements prepared are disarticulated.

Six sauropod genera from four families are positively identified based on remains found at the quarry. The diplodocids are represented by at least three species. Most of the *Diplodocus* remains from the quarry belong to adults, as defined by complete neurocentral fusion. However, they are smaller than all other adult *Diplodocus*. Characters in the DMDQ *Diplodocus* and *Brachiosaurus* vertebrae suggest they may be new species.

DMDQ sauropods with representative diagnostic specimen numbers:

*Apatosaurus* sp. - putative caudal #1 (BYU 5065), rt. fibula (BYU 11420)

*Brachiosaurus* sp. - rt. scapulocoracoid (BYU 9462), sacral complex with three articulated caudal vertebrae (BYU 11434)

*Camarasaurus* sp. - caudal vertebrae (BYU 12852, 12853, 13286), rt. tibia (BYU 17352)


*Diplodocus* sp. - articulated skeleton from presacral 3 to caudal 15+ (BYU 12155), rt. tibia (BYU 12604)

*Haplocanthosaurus* sp. - dorsal vertebrae (BYU 9194, 11506, 17530, 17531, 17689, 17530), rt. tibia (BYU 12865)

*Supersaurus* - caudal vertebra (BYU 9044), holotype scapulocoracoid (BYU 12962)

Additionally, a cervical vertebra (BYU 11617) may belong to *Barosaurus*. If so, this would make a seventh sauropod genus present.

With the recognition that the specimens described herein are referable to *Haplocanthosaurus*, DMDQ is the first – and so far only – known quarry in the world with at least six distinct, valid genera of sauropods present. The quarry is hypothesized to represent a faunal assemblage that was first concentrated and then killed by a severe drought (Richmond and Morris, 1998), so the six sauropod genera might have originally been segregated among different nearby habitats during favorable, non-drought conditions. Still, the presence of at least six sauropod genera within the area that could be concentrated



**Figure 1.** *Haplocanthosaurus* tibiae and dorsal vertebrae

during a severe drought emphasizes the high site-level diversity of Morrison sauropods.

Is DMDQ unique in the Morrison in supporting so many dinosaur taxa? At least nineteen dinosaur genera (six sauropods, eight theropods, and five ornithischians) are currently recognized. Is it a “disaster quarry,” the bone salad resulting from a massive drought drawing all of the surrounding taxa into close quarters? Was it a crossroads site, occupying a migration path, or situated on the boundary between two ecosystems? Evidence from Whitlock et al. indicates this quarry was part of the western region of the Morrison Formation, and DMDQ's sauropod assemblage provides further evidence for the high diversity across the Western and Eastern regions of the Morrison (Whitlock et al., 2018).

## References

- Billbey S. A., Hall, J. E., & Hall D. A. (2000). Preliminary results on a new haplocanthosaurid sauropod dinosaur from the lower Morrison Formation of northeastern Utah. *Journal of Vertebrate Paleontology*, 20(S3), 30A.
- Britt, B. B. (1991). Theropods of Dry Mesa Quarry (Morrison Formation, late Jurassic), Colorado, with emphasis on the osteology of *Torvosaurus tanneri*. *Brigham Young University Geology Studies*, 37, 1-72.
- Foster, J. (2020). *Jurassic West: The dinosaurs of the Morrison Formation and their world* (2nd ed.). Indiana University Press.
- Foster, J. R., Wedel, M. J. (2014). *Haplocanthosaurus* (Saurischia: Sauropoda) from the lower Morrison Formation (Upper Jurassic) near Snowmass, Colorado. *Volumina Jurassica*, 12(2), 197-210.
- Hatcher, J. B. (1903). *Osteology of Haplocanthosaurus, with description of a new species and*

*remarks on the probable habits of the Sauropoda and the age and origin of the Atlantosaurus beds: Additional remarks on Diplodocus*. Vol. 2(1). Carnegie Museum.

McIntosh, J. S., & Williams, M. E. (1988). A new species of sauropod dinosaur, *Haplocanthosaurus delphi* sp. nov., from the Upper Jurassic Morrison Fm. of Colorado. *Kirtlandia*, 43, 3-26.

Richmond, D.R., & Morris, T. H. (1998). Stratigraphy and cataclysmic deposition of the Dry Mesa Dinosaur Quarry, Mesa County, Colorado. *Modern Geology*, 22, 121-143.

Turner, C. E., & Peterson, F. (1999). Biostratigraphy of dinosaurs in the Upper Jurassic Morrison Formation of the Western Interior, USA. *Utah Geological Survey Miscellaneous Publication*, 99, 77-114.

Whitlock, J. A., Trujillo, K. C., & Hanik, G. M. (2018). Assemblage-level structure in Morrison Formation dinosaurs, Western Interior, USA. *Geology of the Intermountain West*, 5, 9-22.

## Technical Session 3: Terrestrial Ecosystems – Triassic – Middle Jurassic (Friday, June 9, 2023, 9:45 AM)

### GEOLOGY AND PALEONTOLOGY OF THE UPPER TRIASSIC CHINLE FORMATION IN THE NORTHERN PARADOX BASIN OF SOUTHEASTERN UTAH: COMPARING SURVEYS FROM CANYONLANDS NATIONAL PARK AND INDIAN CREEK IN BEARS EARS NATIONAL MONUMENT

DeBlieux, Donald D.<sup>1</sup>, Milner, Andrew R.C.<sup>2</sup>, Irmis, Randall B.<sup>3</sup>, Kirkland, James I.<sup>1</sup>